

# IJSTR

*by* Arika Kristiana

---

**Submission date:** 20-Apr-2021 11:29AM (UTC+0700)

**Submission ID:** 1564370909

**File name:** 10\_IJSTR\_arika\_jilid-10-12.pdf (643.12K)

**Word count:** 1673

**Character count:** 7766

# The R-Dynamic Local Irregularity Vertex Coloring Of Graph

A. I. Kristiana, M. I. Utoyo, Dafik, R. Alfarisi, E. Waluyo

**Abstract:** We define the  $r$ -dynamic local irregularity vertex coloring. Suppose  $\lambda : V(G) \rightarrow \{1, 2, \dots, k\}$  is called vertex irregular  $k$ -labeling and  $w : V(G) \rightarrow N$  where  $w(u) = \sum_{v \in N(u)} \lambda(v)$ .  $\lambda$  is called  $r$ -dynamic local irregular vertex coloring, if: (i)  $\text{opt}(\lambda) = \min\{\max\{\lambda_i\}; \lambda_i \text{ vertex irregular } k\text{-labeling}\}$ , (ii) for every  $uv \in E(G)$ ,  $w(u) \neq w(v)$ , and (iii) for every  $v \in V(G)$  such that  $|w(N(v))| \geq \min\{r, d(v)\}$ . The chromatic number  $r$ -dynamic local irregular denoted by  $\chi_{lis}^r(G)$ , is minimum of cardinality  $r$ -dynamic local irregular vertex coloring. We study the  $r$ -dynamic local irregularity vertex coloring of graph and we have found the exact value of chromatic number  $r$ -dynamic local irregularity of some graph.

**Index Terms:**  $r$ -dynamic coloring, local irregularity, vertex coloring.

## 1 INTRODUCTION

GRAPH in this paper are simple and finite. For  $v \in V(G)$ , let  $N(v)$  denote the set of vertices adjacent to  $v$  in  $G$  and  $d(v) = |N(v)|$ . Vertices in  $N(v)$  are neighbors of  $v$ . Montgomery [3] introduced the  $r$ -dynamic coloring. Let  $r$  be a positive integer. An  $r$ -dynamic  $k$ -coloring is a proper vertex  $k$ -coloring such that every vertex  $v$  receives at least  $\min\{r, d(v)\}$ . Furthermore Lai defined  $r$ -dynamic chromatic number that the minimum  $k$ , which  $G$  admits an  $r$ -dynamic  $k$ -coloring and is denoted  $\chi_r(G)$ .

Kristiana, et.al [1] defined local irregularity vertex coloring. Suppose  $l : V(G) \rightarrow \{1, 2, \dots, k\}$  is called vertex irregular  $k$ -labeling and  $w : V(G) \rightarrow N$  where  $w(u) = \sum_{v \in N(u)} l(v)$ ,  $l$  is called local irregularity vertex coloring, if (i)  $\max(l) = \min\{\max\{l_i\}\}$  and (ii) for every  $uv \in E(G)$ ,  $w(u) \neq w(v)$ . Furthermore Kristiana, et.al [2] found chromatic number local irregularity of path graph, cycle graph, complete graph, bipartite complete graph, star graph, and friendship graph. In this paper, we combine  $r$ -dynamic coloring and local irregularity vertex coloring.

## 2 RESULT

In this paper, we present new definition of the  $r$ -dynamic local irregularity vertex coloring of graph and the chromatic number  $r$ -dynamic local irregular. We study the exact value of chromatic number  $r$ -dynamic local irregular of some graphs.

### Definition 1

Let  $\lambda : V(G) \rightarrow \{1, 2, \dots, k\}$  is called vertex irregular  $k$ -labeling and  $w : V(G) \rightarrow N$  where  $w(u) = \sum_{v \in N(u)} \lambda(v)$ .  $\lambda$  is called  $r$ -dynamic local irregular vertex coloring, if:

- $\text{opt}(\lambda) = \min\{\max\{\lambda_i\}; \lambda_i \text{ vertex irregular } k\text{-labeling}\}$
- For every  $uv \in E(G)$ ,  $w(u) \neq w(v)$
- For every  $v \in V(G)$  such that  $|w(N(v))| \geq \min\{r, d(v)\}$ .

### Definition 2

The chromatic number  $r$ -dynamic local irregular denoted by  $\chi_{lis}^r(G)$ , is minimum of cardinality  $r$ -dynamic local irregular vertex coloring.

For  $r = 1$  is called chromatic number local irregular and for  $r = 2$  is called chromatic number dynamic local irregular.




Figure 1. An example of local irregularity  $r$ -dynamic

Illustration of local irregularity  $r$ -dynamic vertex coloring is presented in Figure 1.

### Observation 1

Let be graph  $G$ , where  $N(u) = N(v)$ , graph  $G$  doesn't have local irregularity vertex coloring for  $r \geq 2$ .

Based on Observation 1, some graph don't have local irregularity  $r$ -dynamic for  $r \geq 2$ , namely star graph, path graph with order 3 and bipartite graph.

### Observation 2

Let be  $G$  connected graph, local irregularity  $r$ -dynamic vertex coloring for  $r \geq 2$  have  $\text{opt}(\lambda) \geq 3$ .

### Lemma 1

Graph connected  $G$ ,  $\chi_{lis}^r(G) \geq \chi_{lis}(G)$

Proof: Let  $b : V(G) \rightarrow N$  be local irregularity vertex coloring, for  $uv \in E(G)$ ,  $b(u) \neq b(v)$ .

$$\chi_{lis}(G) = \min\{|b(V(G))|; b \text{ local irregularity vertex coloring}\}$$

Based on Definition 1,  $b$  is vertex irregular  $k$ -labeling such that  $\chi_{lis}(G) \leq |b(V(G))|$ .

Thus,  $\chi_{lis}(G) \leq \min\{|b(V(G))|\} = \chi_{lis}^r(G)$ .  $\square$

- Arika Indah Kristiana, University of Jember, Jember, Indonesia. [arika.fkip@unej.ac.id](mailto:arika.fkip@unej.ac.id)
- Moh. Imam Utoyo, University of Airlangga, Surabaya, Indonesia. [m.i.utoyo@fst.unair.ac.id](mailto:m.i.utoyo@fst.unair.ac.id)
- Dafik, University of Jember, Jember, Indonesia. [d.dafik@unej.ac.id](mailto:d.dafik@unej.ac.id)
- Ridho Alfarisi, University of Jember, Jember, Indonesia. [alfarisi.fkip@unej.ac.id](mailto:alfarisi.fkip@unej.ac.id)
- Eko Waluyo, Islamic Institute of Zainul Hasan, Probolinggo, Indonesia. [ekocasper29@gmail.com](mailto:ekocasper29@gmail.com)

**Theorem 1**

Let  $P_n$  path graph,  $\chi_{lis}^r(P_n) = 4$  where  $n \geq 6$

**Proof:**  $V(P_n) = \{a_i, 1 \leq i \leq n\}$  and  $E(P_n) = \{a_i a_{i+1}, 1 \leq i \leq n-1\}$

Based on Observation 1,  $\text{opt}(\lambda) = 3$  and Based on Lemma 1, the lower bound chromatic number r-dynamic is  $\chi_{lis}^r(P_n) \geq \chi_{lis}^r(P)$ . Further, it will be shown the upper bound, we define  $\lambda : V(P_n) \rightarrow \{1, 2, 3\}$ .

Case 1. For  $n \equiv 0 \pmod 3$

$$\lambda(a_i) = \begin{cases} 1, & i \equiv 1 \pmod 3, 1 \leq i \leq n \\ 2, & i \equiv 2 \pmod 3, 1 \leq i \leq n \\ 3, & i \equiv 0 \pmod 3, 1 \leq i \leq n \end{cases}$$

It easy to see  $\text{opt}(\lambda) = 3$  and weight function as follows:

$$w(a_i) = \begin{cases} 2, & i = 1, n \\ 3, & i \equiv 0 \pmod 3, 2 \leq i \leq n-1 \\ 4, & i \equiv 2 \pmod 3, 2 \leq i \leq n-1 \\ 5, & i \equiv 1 \pmod 3, 2 \leq i \leq n-1 \end{cases}$$

Case 2. For  $n \equiv 1 \pmod 3$

$$\lambda(a_i) = \begin{cases} 1, & i \equiv 1 \pmod 3, 1 \leq i \leq n-2 \\ 2, & i = n-1, n \text{ or } i \equiv 2 \pmod 3, 1 \leq i \leq n-2 \\ 3, & i \equiv 0 \pmod 3, 1 \leq i \leq n-2 \end{cases}$$

It easy to see  $\text{opt}(\lambda) = 3$  and weight function as follows:

$$w(a_i) = \begin{cases} 2, & i = 1, n \\ 3, & i = n-2 \text{ or } i \equiv 0 \pmod 3, 2 \leq i \leq n-2 \\ 4, & i = n-1 \text{ or } i \equiv 2 \pmod 3, 2 \leq i \leq n-2 \\ 5, & i \equiv 1 \pmod 3, 2 \leq i \leq n-2 \end{cases}$$

Case 3. For  $n \equiv 2 \pmod 3$

$$\lambda(a_i) = \begin{cases} 1, & i = n \text{ or } i \equiv 0 \pmod 3, 1 \leq i \leq n-1 \\ 2, & i \equiv 2 \pmod 3, 1 \leq i \leq n-1 \\ 3, & i \equiv 1 \pmod 3, 1 \leq i \leq n-1 \end{cases}$$

It easy to see  $\text{opt}(\lambda) = 3$  and weight function as follows:

$$w(a_i) = \begin{cases} 2, & i = 1, n-1 \\ 3, & i = n \text{ or } i \equiv 1 \pmod 3, 2 \leq i \leq n-1 \\ 4, & i \equiv 2 \pmod 3, 2 \leq i \leq n-1 \\ 5, & i \equiv 0 \pmod 3, 2 \leq i \leq n-1 \end{cases}$$

For every  $uv \in E(P_n)$ ,  $u = a_i, v = a_{i+1}, 1 \leq i \leq n-1$  obtained  $w(a_i) \neq w(a_{i+1})$ . For  $a_i \in V(P_n)$  such that  $|w(N(a_i))| \geq \min\{r, d(a_i)\}$ . Based on Definition 1,  $w$  is called local irregularity r-dynamic. Weight function obtain  $|w(V(P_n))| = 4$ . Thus,  $\chi_{lis}^r(P_n) \leq 4$ .

Hence,  $\chi_{lis}^r(P_n) = 4$ . The proof is complete.  $\square$

Illustration the r-dynamic local irregularity vertex coloring of path graph is presented in Figure 2.




Figure 2. The 2-dynamic local irregularity of  $P_9$

**Theorem 2**

Let  $C_n$  be cycle graph, for  $n \geq 5$

$$\chi_{lis}^r(C_n) = \begin{cases} 3, & n \equiv 0 \pmod 3 \\ 4, & n = 7 \\ 5, & n \equiv 1, 2 \pmod 3, n \neq 7 \end{cases}$$

**Proof:**  $V(C_n) = \{a_i, 1 \leq i \leq n-1\}$  and  $E(C_n) = \{a_i a_{i+1}, 1 \leq i \leq n-1\} \cup \{a_n a_1\}$ . Based on Observation 1,  $\text{opt}(\lambda) = 3$  and Based on Lemma 1, the lower bound chromatic number r-dynamic is  $\chi_{lis}^r(C_n) \geq$

$\chi_{lis}^r(C_n) = 3$ . Further, it will be shown the upper bound, we define  $\lambda : V(C_n) \rightarrow \{1, 2, 3\}$ .

Case 1. For  $n \equiv 0 \pmod 3$

$$\lambda(a_i) = \begin{cases} 1, & i \equiv 1 \pmod 3, 1 \leq i \leq n \\ 2, & i \equiv 2 \pmod 3, 1 \leq i \leq n \\ 3, & i \equiv 0 \pmod 3, 1 \leq i \leq n \end{cases}$$

It easy to see  $\text{opt}(\lambda) = 3$  and weight function as follows:

$$w(a_i) = \begin{cases} 3, & i \equiv 0 \pmod 3, 1 \leq i \leq n \\ 4, & i \equiv 2 \pmod 3, 1 \leq i \leq n \\ 5, & i \equiv 1 \pmod 3, 1 \leq i \leq n \end{cases}$$

For every  $uv \in E(C_n)$ ,  $u = a_i, v = a_{i+1}, 1 \leq i \leq n-1$  obtained  $w(a_i) \neq w(a_{i+1})$  and  $u = a_n, v = a_1$  obtained  $w(a_n) \neq w(a_1)$ . For  $a_i \in V(C_n)$  such that  $|w(N(a_i))| \geq \min\{r, 2\}$ . Based on Definition 1,  $w$  is called local irregularity r-dynamic. Weight function obtain  $|w(V(C_n))| = 3$ . Thus,  $\chi_{lis}^r(C_n) \leq 3$ .

Hence  $\chi_{lis}^r(C_n) = 3$

Case 2. For  $n = 7$

$$\lambda(a_i) = \begin{cases} 1, & n = 2, 4 \\ 2, & n = 5, 6, 7 \\ 3, & n = 1, 3 \end{cases}$$

It easy to see  $\text{opt}(\lambda) = 3$  and weight function as follows:

$$w(a_i) = \begin{cases} 2, & i = 3 \\ 3, & i = 1, 5 \\ 4, & i = 2, 6 \\ 5, & i = 4, 7 \end{cases}$$

For every  $uv \in E(C_n)$ ,  $u = a_i, v = a_{i+1}, 1 \leq i \leq n-1$  obtained  $w(a_i) \neq w(a_{i+1})$  and  $u = a_n, v = a_1$  obtained  $w(a_n) \neq w(a_1)$ . For  $a_i \in V(C_7)$  such that  $|w(N(a_i))| \geq \min\{r, 2\}$ . Based on Definition 1,  $w$  is called local irregularity r-dynamic. Weight function obtain  $|w(V(C_n))| = 4$ . Thus,  $\chi_{lis}^r(C_n) \leq 4$ .

Hence  $\chi_{lis}^r(C_7) = 4$

Case 3. For  $n \equiv 1, 2 \pmod 3, n \neq 7$

Subcase 1. For  $n \equiv 1 \pmod 3, n \neq 7$

$$\lambda(a_i) = \begin{cases} 1, & i = 2, n-1 \text{ or } i \equiv 1 \pmod 3, 4 \leq i \leq n-3 \\ 2, & i = n \text{ or } i \equiv 2 \pmod 3, 4 \leq i \leq n-3 \\ 3, & i = 1, n-2 \text{ or } i \equiv 0 \pmod 3, 3 \leq i \leq n-3 \end{cases}$$

It easy to see  $\text{opt}(\lambda) = 3$  and weight function as follows:

$$w(a_i) = \begin{cases} 2, & i = 3, n-2 \\ 3, & i = 1 \text{ or } i \equiv 0 \pmod 3, 4 \leq i \leq n-4 \\ 4, & i = n \text{ or } i \equiv 2 \pmod 3, 4 \leq i \leq n-4 \\ 5, & i = n-1 \text{ or } i \equiv 2 \pmod 3, 4 \leq i \leq n-4 \\ 6, & i = 2, n-3 \end{cases}$$

Subcase 2. For  $n \equiv 2 \pmod 3$

$$\lambda(a_i) = \begin{cases} 1, & i \equiv 1 \pmod 3, 1 \leq i \leq n-1 \\ 2, & i \equiv 2 \pmod 3, 1 \leq i \leq n-1 \\ 3, & i = n \text{ or } i \equiv 0 \pmod 3, 1 \leq i \leq n-1 \end{cases}$$

It easy to see  $\text{opt}(\lambda) = 3$  and weight function as follows:

$$w(a_i) = \begin{cases} 2, & i = n \\ 3, & i \equiv 0 \pmod 3, 1 \leq i \leq n-2 \\ 4, & i \equiv 2 \pmod 3, 1 \leq i \leq n-2 \\ 5, & i \equiv 1 \pmod 3, 1 \leq i \leq n-2 \\ 6, & i = n-1 \end{cases}$$

For every  $uv \in E(C_n)$ ,  $u = a_i, v = a_{i+1}, 1 \leq i \leq n-1$  obtained  $w(a_i) \neq w(a_{i+1})$  and  $u = a_n, v = a_1$  obtained  $w(a_n) \neq w(a_1)$ . For  $a_i \in V(C_n)$  such that  $|w(N(a_i))| \geq \min\{r, 2\}$ . Based on Definition 1,  $w$  is

called local irregularity  $r$ -dynamic. Weight function obtain  $|w(V(C_n))| = 5$ . Thus,  $\chi_{lis}^r(C_n) \leq 5$ . Hence  $\chi_{lis}^r(C_n) = 5$ . The proof is complete.  $\square$

Illustration the  $r$ -dynamic local irregularity vertex coloring of cycle graph is presented in Figure 3.




Figure 3. The 2-dynamic local irregularity of  $C_7$

### Theorem 3

Let  $K_n$  be complete graph,  $\chi_{lis}^r(K_n) = n$

Proof:  $V(K_n) = \{a_i, 1 \leq i \leq n\}$ . Suppose  $u, v \in V(K_n)$ , Based on Observation 3,  $N(u) - \{v\} = N(v) - \{u\}$  so that  $\lambda(u) \neq \lambda(v)$ . It show labeling of every vertex in complete graph as different. So  $\text{opt}(\lambda) = n$ . Based on Lemma 1,  $\chi_{lis}^r(K_n) \geq \chi_{lis}(K_n) = n$ . Further, to show the upper bound, we define  $\lambda : V(K_n) \rightarrow \{1, 2, \dots, n\}$  where  $\lambda(a_i) = i, 1 \leq i \leq n$ . weight function is  $w(a_i) = \frac{n(n+1)}{2} - i$ . Because  $i = 1, 2, \dots, n$ , where  $|w(V(K_n))| = n$  so that  $n = \chi_{lis}(K_n) \leq \chi_{lis}^r(K_n) \leq |w(V(K_n))| = n$ . Thus,  $\chi_{lis}^r(K_n) = n$ . The proof is complete.  $\square$

### 3 CONCLUSION

In this paper we have studied the  $r$ -dynamic local irregularity vertex coloring. We have concluded the exact value of the chromatic number  $r$ -dynamic local irregular of some graphs, namely path graph, cycle graph and complete graph.

### 3 ACKNOWLEDGMENT

We gratefully acknowledgment the support from University of Jember of 2019.

### REFERENCES

- [1] M. Alishahi, "on the dynamic coloring of graphs," *Discrete Applied Mathematics*, vol. 159, pp. 152-156, 2011.
- [2] S. Jahanbekam, J. Kim, Suil. O and D. B. West, "On  $r$ -dynamic coloring of graphs," *Discrete Applied Mathematics*, vol. 206, pp. 65-72, 2016.
- [3] A.I. Kristiana, M. I. Utoyo, Dafik, Slamin, R. Alfarisi, and I. H. Agustin, "Local irregularity vertex coloring of graph," *International Journal of Civil Engineering and Technology*, vol. 10, issue 3, pp. 1606-1616. 2019.
- [4] A. I. Kristiana, M. I. Utoyo, Dafik, R. Alfarisi, I. H. Agustin, and E. Waluyo., " On the chromatic number local irregularity of related wheel graph," *Journal of Physics: Conf. Series*, 1211, 012003, 2019.

- [5] B. Montgomery, "Dynamic coloring of graphs," PhD dissertation, Department of Mathematics West Virginia University, Morgantown, 2001.
- [6] A. Taherkhani, "On  $r$ -dynamic chromatic number of graphs," *Discrete Applied Mathematics*, vol. 201, pp. 222-227, 2016.

16%

SIMILARITY INDEX

7%

INTERNET SOURCES

15%

PUBLICATIONS

%

STUDENT PAPERS

## PRIMARY SOURCES

1	Jia Bao Liu, Sana Akram, Muhammad Javaid, Abdul Raheem, Roslan Hasni. " Bounds of Degree-Based Molecular Descriptors for Generalized -sum Graphs ", Discrete Dynamics in Nature and Society, 2021 Publication	4%
2	<a href="http://www.revistaproyecciones.cl">www.revistaproyecciones.cl</a> Internet Source	2%
3	Ridho Alfarisi, Dafik, Arika Indah Kristiana. "Resolving domination number of graphs", Discrete Mathematics, Algorithms and Applications, 2019 Publication	2%
4	<a href="http://www.ijstr.org">www.ijstr.org</a> Internet Source	2%
5	Sogol Jahanbekam, Jaehoon Kim, Suil O, Douglas B. West. "On r-dynamic coloring of graphs", Discrete Applied Mathematics, 2016 Publication	1%
6	<a href="http://penelitian.uisu.ac.id">penelitian.uisu.ac.id</a> Internet Source	1%

---

7 Dian Puspita Hapsari, Imam Utoyo, Santi Wulan Purnami. "Fractional Gradient Descent Optimizer for Linear Classifier Support Vector Machine", 2020 Third International Conference on Vocational Education and Electrical Engineering (ICVEE), 2020 1 %

Publication

---

8 Ali Taherkhani. " On -dynamic chromatic number of graphs ", Discrete Applied Mathematics, 2016 1 %

Publication

---

9 "Co-Secure Set Domination in Graphs", International Journal of Recent Technology and Engineering, 2020 1 %

Publication

---

10 Dan Yi, Junlei Zhu, Lixia Feng, Jiaxin Wang, Mengyini Yang. "Optimal r-dynamic coloring of sparse graphs", Journal of Combinatorial Optimization, 2019 1 %

Publication

---

11 R. S. Booth. "Location of Zeros of Derivatives. II", SIAM Journal on Applied Mathematics, 1969 1 %

Publication

---

Exclude bibliography  On